



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

That primes are quite irregularly distributed, will be plain from this:

The 26,379th hundred contains no prime.

The 27,050th hundred contains 17 primes.

Even when they are taken in large masses, the irregularity is still apparent, thus:—

The 27th hundred thousand contains 6,762 primes.

The 28th hundred thousand contains 6,714 primes.

The 29th hundred thousand contains 6,744 primes.

For the numerical data here given I am indebted to Gauss, (Vol. above quoted.)  
G. W. HILL.

$$\text{The formula } N = \frac{x}{A \log. x - B}$$

referred to by Prof. Brooks in No. 3, requires that the logarithm should be hyperbolic, which he neglects to state.

This formula is fully and ably discussed in Vol. II, § VIII, p. 65 of *Legendre's Theorie des Nombres*, 3d Edition, 1830.

The formula adopted by him is

$$N = \frac{x}{h. \log. x - 1.08366}$$

Many years since I became interested in this matter, and I then used the formula

$$N = \frac{x}{1.001 h. \log. x - 1.09}$$

G. R. P.

### SOLUTIONS OF PROBLEMS IN NO. 2.

Solutions have been received as follows: R. J. Adcock solved 7; Irving P. Church solved 6; G. W. Hill solved 9; Prof. A. Hall solved 10; H. Heaton solved 6, 7, 8, 9 and 10; Prof. D. Kirkwood solved 7; Artemas Martin solved 5, 6 and 7; Prof. J. Scheffer solved 5 and 7; James Stott solved 6; Walter Siverly solved 8; and Prof. W. Wylie solved 7. David Wickersham and William Hoover sent solutions to Problems in No. 1 too late for notice in No. 3.

5. "In a plane triangle there are given the three lines bisecting the angles,  $a$ ,  $b$  and  $c$ , to find the sides."

SOLUTION BY PROF. J. SCHEFFER, COLLEGE OF ST. JAMES, MD.

Let  $CF = a$ ,  $BE = b$ ,  $AD = c$  be the three bisecting lines, and let